

Danger Signs

If a client/child has any of these signs he or she might die and needs to be taken immediately to the nearest health facility

Danger Signs
Convulsions or fits now or within the past 2 days
Coma / Loss of consciousness
Client/child is confused or very sleepy - cannot be woken
Extreme weakness – unable to stand or sit without support
Very Hot – with temperature of 38.5 OC or more
Very Cold –with temperature of 35.0 OC or less
Vomiting everything - cannot keep down food or drink
Not able to drink or breastfeed
Severe anaemia – very pale palms, fingernails, eyelids
Yellow eyes
Difficulty in breathing
Severe dehydration - sunken eyes, sunken fontanelle, skin pinch, coated tongue

What to do if a client/child has any of these danger signs:

1. Check blood using Quick Malaria Test
2. If test is: **POSITIVE** - Treat client/child with Antimalarial Suppository (Rectal Artesunate)
NEGATIVE - Do not give any medicine; refer to health unit for treatment instead
3. Write Emergency Referral Form – use **RED FORM**
4. Instruct caretaker to take client/child to the nearest health unit immediately without delay

Any illness in a young baby less than 2 months old:

A young baby can quickly become dangerously sick. Refer baby without delay to the nearest health facility – use **RED FORM**

DO NOT GIVE RECTAL ARTESUNATE or any other medicine as this may not be safe in a baby